

Page 1.

NAME - M. Sohrab

ID - 14589.

Submitted to <sup>Sir</sup> RaZa Ahmed Khan

Subject Basic Statistic.

Q1:-  
(a)

$x$	$f(x)$	$f(x)$	$\frac{1}{x}$	$f(\frac{1}{x})$	$\log(x)$	$f(\log x)$
1	4	4	1.0	4	0	0
2	13	26	0.50	6.5	0.3	3.9
3	9	27	0.33	2.97	0.477	4.29
4	4	16	0.25	1	0.6	2.4
5	1	5	0.2	0.2	0.69	0.69
		<u>78</u>		<u>14.67</u>		<u>11.28</u>

$$A.M = \frac{\sum f(x)}{\sum f} = \frac{78}{31} = \boxed{2.5161}$$

$$G.M = \text{Antilog} \left[ \frac{\sum f(\log x)}{\sum f} \right] = \text{Antilog} \left[ \frac{11.28}{31} \right] \\ = \boxed{2.311}$$

$$H.M = \frac{\sum f}{\sum f(\frac{1}{x})} = \frac{31}{14.67} = \boxed{2.11}$$

(2)

(b) A.M

Marks	f	Mid Point (u)	fu
0-9	2	4.5	9
10-19	31	14.5	449.5
20-29	73	24.5	1788.5
30-39	85	34.5	2932.5
40-49	28	44.5	1246
	$\Sigma f = 219$		$\Sigma fu = 6425.5$

$$A.M = \frac{\Sigma fu}{\Sigma f} = \frac{6425.5}{219}$$

(b) G.M

$$= 29.3401$$

Marks	f	u	fu	$f \times \log(u)$
0-9	2	4.5	9	1.3084
10-19	31	14.5	449.5	36.0024
20-29	73	24.5	1788.5	101.4091
30-39	85	34.5	2932.5	130.7146
40-49	28	44.5	1246	46.1540
	$\Sigma f = 219$		$\Sigma fu = 6425.5$	$\Sigma f \times \log u = 315.5865$

$$G.M = \text{Anti} \log \left( \frac{\Sigma f \log(u)}{\Sigma f} \right) = \frac{315.5865}{219}$$

$$= 1.4410$$

$$= 27.6057$$

(15)  
(3)

H.M

Marks	f	n	fu	$\frac{fu}{n}$
0-9	2	4.5	9	0.444
10-19	31	14.5	449.5	2.1379
20-29	73	24.5	1788.5	2.9795
30-39	85	34.5	2932.5	2.4637
40-49	28	44.5	1246	0.6292
	<u><math>\Sigma f = 219</math></u>		<u><math>\Sigma fu = 6425</math></u> 5	<u>8.6547</u>

$$H.M = \frac{\Sigma f}{\Sigma \frac{fu}{n}} = \frac{219}{8.6547} = 25.3041.$$

(5)  
(4)

Q2:- Part a

n	f	C.F.
1	4	4
2	13	17
3	9	26
4	4	30
5	1	31

$$\text{Median } \left[ \frac{n+1}{2} \right]^{\text{th}} = \frac{32}{2} = 16^{\text{th}}$$

$$\boxed{\text{Median} = 2}$$

$\boxed{\text{Mode} = 2}$  Because it has highest frequency.

(5)

Q2:-(b)

Marks	C.B	f	C.F
0-9	-0.5-9.5	2	2
10-19	9.5-19.5	31	33
20-29	19.5-29.5	73	106
30-39	29.5-39.5	85	191
40-49	39.5-49.5	28	219
		<u>219</u>	

$$\Rightarrow \text{Median} = l + \frac{N/2 - m}{f} + c$$

$$\begin{aligned} \therefore l &= 29.5, N = 219 \\ m &= 106, c = 10 \\ f &= 85 \end{aligned}$$

$$\text{Median} = 29.5 + \frac{(219/2) - 106 \times 10}{85}$$

$$\Rightarrow \text{Median} = 29.5 + \frac{(219/2) - 106 \times 10}{85}$$

$$\boxed{\text{Median} = 29.91 = 30}$$

$$\Rightarrow \text{Mode} = l + h \left[ \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right]$$

$$= 29.5 + 10 \left[ \frac{85 - 73}{2(85) - 73 - 28} \right]$$

$$\boxed{\text{Mode} = 31.23}$$

(6)

Q30- (a)

Marks	C.B	f	C.F
0-9	-0.5-9.5	2	2
10-19	9.5-19.5	31	33
20-29	19.5-29.5	71	106
30-39	29.5-39.5	85	191
40-49	39.5-49.5	28	219

$$Q_1 \text{ class} = \frac{N}{4} = \frac{219}{4} = 54.75$$

$$Q_3 \text{ class} = 3\left[\frac{N}{4}\right] = 3 \times 54.75 = 164.25$$

$$Q_1 = \frac{L + \frac{N}{4} - m_1 \times c}{f_1}$$

$$= \frac{19.5 + 54.75 - 33 \times 10}{71} \quad \boxed{Q_1 = 22.56}$$

$$Q_3 = \frac{L_3 + 3\left(\frac{N}{4}\right) - m_3 \times 10}{f_3}$$

$$= \frac{29.5 + 164.25 - 106 \times 10}{85} \quad \boxed{Q_3 = 36.35}$$

$$IQR = 36.35 - 22.56 = 13.79$$

$$\text{Semi IQR} = \frac{13.79}{2} = 6.895$$

Q3:-(b)

C.B	$x$	$f$	$\bar{x}$	$x - \bar{x}$	$(x - \bar{x})^2$
0.5-9.5	4.5	2	29.33	-24.83	616.52
9.5-19.5	14.5	31	29.33	-14.83	219.92
19.5-29.5	24.5	73	29.33	-4.83	23.32
29.5-39.5	34.5	85	29.33	5.17	26.73
39.5-49.5	44.5	28	29.33	15.17	230.12
		<u>219</u>			

$$\text{variance} = \sigma^2 = \frac{\sum f(x - \bar{x})^2}{\sum f}$$

$$\sigma^2 = \frac{18468.33}{219}$$

$$\sigma^2 = 84.33$$

$$\begin{array}{r} \sum f(x - \bar{x})^2 \\ 1233.04 \\ 6817.52 \\ 1702.36 \\ 2272.05 \\ 6443.38 \\ \hline 18468.33 \end{array}$$

$$\text{Standard deviation} = \sigma = \sqrt{\sigma^2}$$

$$= \sqrt{84.33}$$

$$\Rightarrow \sigma = 9.18$$



Sign in to edit and save changes to this...



**Q4)**

**A)Range & Quartile Range & Semi Inter Quartile Range**

**Ans)**

**Range:** The **Range is the** difference between the lowest and highest values.

**Example:** In {4, 6, 9, 3, 7} the lowest value is 3, and the highest is 9. So the **range** is  $9 - 3 = 6$ .

**Quartile Range:** The interquartile **range (IQR)** is a measure of variability, based on dividing a data set into **quartiles**. **Quartiles** divide a rank-ordered data set into four equal parts. The values that divide each part are called the first, second, and third **quartiles**; and they are denoted by Q1, Q2, and Q3, respectively.

**Example:**

5, 7, 4, 4, 6, 2, 8

Put them in order: 2, 4, 4, 5, 6, 7, 8. Cut the list into quarters: And the result is: **Quartile 1 (Q1) = 4. Quartile 2 (Q2)**, which is also the Median, = 5.

**Semi inter quartile range:**

The semi-interquartile range is a measure of spread or dispersion. It is computed as one half the difference between the 75th percentile [often called (Q3)] and the 25th percentile (Q1). The formula for semi-interquartile range is therefore:  $(Q3-Q1)/2$ .

The semi-interquartile range is half of the difference between the upper quartile and the lower quartile. In the previous example, the quartiles were Q 1 = 4 and. Q 3 = 11 . The semi-interquartile range is:  $1/2 ( Q 3 - Q 1 )$





## Document (15) - Read-only



Sign in to edit and save changes to this... 

**Example:**

**Question:** Find the Quartile Deviation for the following set of data:

{490, 540, 590, 600, 620, 650, 680, 770, 830, 840, 890, 900}

**Step 1:** Find the **first** quartile,  $Q_1$ .

This is the median of the lower half of the set {490, 540, 590, 600, 620, 650}.

$$Q_1 = (590 + 600) / 2 = 595.$$

**Step 2:** Find the **third** quartile,  $Q_3$ .

This is the median of the upper half of the set {680, 770, 830, 840, 890, 900}.

$$Q_3 = (830 + 840) / 2 = 835.$$

**Step 3:** Subtract Step 1 from Step 2.

$$835 - 595 = 240.$$

**Step 4:** Divide by 2.  $240 / 2 = 120$

**QB) Variance & Standard Deviation & Coefficient of variation?****Ans**

**Variance:** Variance ( $\sigma^2$ ) in statistics is a measurement of the spread between numbers in a data set. That is, it measures how far each number in the set is from the mean and therefore from every other number in the set.

**Example:**

for the numbers 1, 2, and 3 the mean is 2 and the **variance** is 0.667.

**Standard Deviation:** The **standard deviation** is a statistic that measures the dispersion of a dataset relative to its mean and is calculated as the square root of the variance. The **standard deviation** is calculated as the square root of variance by determining each data point's **deviation** relative to the mean.

**Example:** 5, 15, 15, 14, 16 and 2, 7, 14, 22, 30. However, the second is clearly more spread out.

**Coefficient of variation:**

Sign in to edit and save changes to this...



The coefficient of variation (relative standard deviation) is a statistical measure of the dispersion of data points around the mean. The metric is commonly used to compare the data dispersion between distinct series of data. Unlike the standard deviation that must always be considered in the context of the mean of the data, the coefficient of variation provides a relatively simple and quick tool to compare different data series.

### **Example:**

Fred wants to find a new investment for his portfolio. He is looking for a safe investment that provides stable returns. He considers the following options for investment:

- **Stocks:** Fred was offered stock of ABC Corp. It is a mature company with strong operational and financial performance. The volatility of the stock is 10% and the expected return is 14%.
- **ETFs:** Another option is an Exchange-Traded Fund (ETF) which tracks the performance of the S&P 500 index. The ETF offers an expected return of 13% with a volatility of 7%.
- **Bonds:** Bonds with excellent credit ratings offer an expected return of 3% with 2% volatility.

