

Final-Term Assignment

Course Title: **Human Physiology II**

Rad 2nd semester section A

Instructor: Dr. M .Shahzeb khan (PT)

Marks: 50

Note:

- **Attempt all questions, all questions carry equal marks.**
- **Answer Briefly and to the point, avoid un-necessary details**

Q1: (A) How stimulus of smell moves from nostril to brain? Make a Diagram as well

Q 2: (A) What is difference between Haemostasis, Haematopoiesis and Homeostasis?

(B) What is Erythroblastosis fetalis?

Q3: (A) What is Immunity? Explain different types of immunity (B)

What is difference between Antigen and Antibody?

Q4: (A) Write down different functions of Antibody

(B) Write difference between Primary and secondary response to an antigen

Q5: Write difference between cell mediated and Antibody Mediated Immunity

Physiology

Semester 2nd

Program B.S Radiology

Section A

Name Tahsin Kamal

ID. 16131

QNo (3)A

Ans

Immunity

The State of being resistant to reinfection with a pathogen . the State of protection against forging pathogen or substance (antigens)

Different types of immunity Innate immunity

We are all Born with some level of immunity to Invaders

Adaptive (acquired immunity

This protect from pathogen develop as we go through life

Passive immunity

This type of immunity is borrowed from another source but it does not last

Indefinitely

Immunization

QNo (3)B

Ans

Antigens

Antigens are substances that trigger the body to cause an immune response. The body perceives antigens as harmful substances and it does its best to eliminate them by producing antibodies.

What antigens do is they activate lymphocytes (white blood cells) responsible for fighting off infection.

Antibodies are also known as immunoglobulins. They are proteins that have a distinct Y shape and are produced by B cells as a result of exposure to antigens.

The antibody has paratope responsible for recognizing a specific epitope of antigen. It is the one responsible for antibody's lock and key binding action. What antibodies do is they get rid of antigens from the body.

QNo (1)

Ans . The ability to smell comes from specialized sensory cells, called olfactory sensory neurotransmitter

.found in a small patch of tissue high inside the nose.

.These cells connect directly to the brain.

. Each olfactory neuron has one odor receptor.

.Microscopic molecules released by substances around us— whether it's coffee brewing t—stimulate these receptors.

. once the neurons detect the molecules, they send messages to your brain, which identifies the smell. There are more smells in the environment than there are receptors, and any given molecule may stimulate a combination of receptors, creating a unique representation in the brain.

.Smells reach the olfactory sensory neurons through two pathways.

.The first pathway is through your nostrils.

. The second pathway is through a channel that connects the roof of the throat to the nose. **QN.2(A)**

Ans. QN.2

Ans. Haemostasis: is the of wound healing. This involves blood clotting.

Haemostasis has three major steps: 1) vasoconstriction, 2) temporary blockage of a break by a platelet plug, and 3) blood coagulation, or formation of a fibrin clot.

These processes seal the hole until tissues are repaired.

Haematopoiesis is the process of forming blood cells, which occurs during embryogenesis and throughout life.

Homoestasis is the process by which variables are regulated so that internal conditions remain stable and relatively constant. Examples of homeostasis include the regulation of temperature and the balance between acidity and alkalinity (pH). It is a process that maintains the stability of the human body's internal environment in response to changes in external conditions.

QN2.(B)

Ans. **Erythroblastosis fetalis** is a severe medical condition that most commonly results from incompatibility between certain blood types of a woman who is pregnant and the fetus. The condition involves a component of blood called Rh factor. Rh factor is an inherited protein, found on the surface of red blood cells. Not everyone has this protein.

QN.4(A)

Ans. Functions of Antibody

1. IgG provides long term protection because it persists for months and years after the presence of the antigen that has triggered their production.

IgG protect against bacteria, viruses, neutralise bacterial toxins, trigger complement protein systems and bind antigens to enhance the effectiveness of phagocytosis.

Main function of IgA is to bind antigens on microbes before they invade tissues. It aggregates the antigens and keeps them in the secretions so when the secretion is expelled, so is the antigen.

IgA are also first defense for mucosal surfaces such as the intestines, nose, and lungs.

IgM is involved in the ABO blood group antigens on the surface of RBCs.

IgM enhance ingestions of cells by phagocytosis.

IgE bind to mast cells and basophils which participate in the immune **response**.

QN.4(B)

Ans. Primary response to an antigen

- 1.This occurs as a result of primary contact with an antigen
- 2.Responding cell is naïve B-cell and T-cell. .
- 3.Lag phase is often longer (4-7 days), sometimes as long as weeks or months
- 4.Level of antibody reaches peak in 7 to 10 days.
- 5.It takes longer time to establish immunity.
- 6.First antibody produced is mainly IgM.
- 8.Antibody level declines rapidly.
- 9.Affinity of antibody is lower for its antigen

Secondary response to an antigen

- 1.This occurs as a result of second and subsequent exposure of the same antigen
- 2.Responding cell is memory cell.
- 3.Lag phase is shorter (1-4 days) due to the presence of memory cell.
- 4.Level of antibody reaches peak in 3 to 5 days.
- 5.Takes shorter time to establish immunity.
- 6.Usually 100-1000 times more antibodies are produced.
- 8.Antibody level remain high for longer period.

QNo (5)

Antibodies mediated immunity's

- 1.Definition.The immunity mediated by macromolecules found in the extracellular body fluids is called humoral immunity. (“humor” a medieval term for body fluid)
- 2.Mediator.The main cell involved in humoral immunity are B-cells.
- 3.Components.B cells, T cells, and macrophages. .

