DPT 4th Course Title: Pharmacology I

Student Name: Syed Muhammad Raahim

Student ID: 14838

Note:

- Paper is divided into two questions, Q1 includes 15 MCQs and Q2 includes 15 True/False statements
- Each MCQ or T/F carry one mark with grand total of 30 marks
- **Highlight** or <u>underline</u> the appropriate option
- Before marking, read every statement carefully to understand the actual sense of question

Q1. Select and highlight the most appropriate option for the following questions

- 1. Which of the following statement is not true
 - a. Oral administration has lower chances of systemic infection
 - b. Activated charcoal easily advert the toxic effects occurred through oral route
 - c. Oral administration has comparatively good absorption
 - d. No specially trained personnel is required for oral administration
- 2. General anesthesia is characterized by _____
 - a. Revocable loss of senses and conscious
 - b. Temporary loss of memory
 - c. Reduced muscle contraction
 - d. All of the above
- 3. Which one of the following routes provides long term release of drugs/medicine
 - a. Subcutaneous
 - b. Intravascular
 - c. Sublingual
 - d. Intramuscular
- 4. All of given statements are correct, except
 - a. Active diffusion is the most common mechanism of absorption

- b. Carrier saturation is always involved in active diffusion
- c. Active transportation involves the utility of adenosine triphosphate

d. Specific proteins are require for carrying drug to the target

- 5. Bioavailability of insulin is negligible through oral administration because of
 - a. Instability in gastric pH
 - b. Degradation by stomach enzymes
 - c. First-pass hepatic metabolism
 - d. All of the above
- 6. Which stage of anesthesia is the most desired for any procedure
 - a. Stage I
 - b. Stage II
 - c. Stage III
 - d. Stage IV
- 7. Due to tight junction of capillaries in brain, only those drugs can be moved that are
 - a. Facilitated by carriers
 - b. Appropriately charged
 - c. Hydrophobic in nature
 - d. Both b. and c.
 - e. Both a. and c. '
- 8. In a healthy adult individual, aminoglycosides; with properties of low molecular weight and lyophobic nature, has the distribution of almost

<mark>a. 20%</mark>

- b. 6%
- c. 60%
- d. None of the above

- When the plasma concentration of drug reaches to 100 after 70, metabolism of drug turns from ______ to _____.
 - a. First order kinetics to zero order kinetics
 - b. Non-linear kinetics to linear kinetics
 - c. No effect on metabolism
 - d. Both a. and b.
- 10. ______ receptors transduces signals from release of serotonin
 - a. Enzymatic-linked receptors
 - b. Ligand-gated receptors
 - c. G-protein coupled receptors
 - d. Intracellular receptors
- 11. By altering the dose of ______ in graded manner, it can be used as anesthetic, anxiolytic and sedative
 - a. Lidocaine
 - b. Diazepam
 - c. Phenytoin
 - d. Both a. and b.
 - e. None of the above
- 12. All the statements are correct, except
 - a. Hierarchical neuronal system has fast conducting fibers, because of neuron myelination
 - b. Functionally metabotropic receptor belongs to diffused neuronal system
 - c. Drugs that pharmacologically affects hierarchical/diffused system have greater effect on responsiveness and hunger etc.
 - d. None of the above
- 13. Therapeutic effect of morphine is mediated through activation of _____
 - a. Endorphin, dynorphin etc.

- b. GABA receptors
- c. Serotonin receptors
- d. Both a. and b.
- e. None of the above
- 14. Depending on subtype, all neurotransmitters can produce both excitation and inhibition of neuron, Except
 - a. Serotonin
 - b. Dopamine
 - c. Glutamic acid
 - d. None of the above
- 15. Which of the following drug have impact on patient weight
 - a. Phenobarbitone
 - b. Lorazepam
 - c. Valproic acid
 - d. Oxcarbazepine

Q2: For the following questions, highlight/underline True or False.

- 1. Respiratory and cardiovascular support must provide at anesthesia i.e. stage of medullary paralysis (True/False)
- 2. Hydrophobic drugs usually cannot move through passive diffusion (True/False)
- 3. If the bond length between drug and receptor is small than it will lead to strengthening of their interaction (True/False)
- 4. Drug solubility increases in diarrheal condition due to increase water content, hence increasing the absorption (True/False)
- Generally, neuronal excitatory signal are transduced by opening of interconnected ion channels i.e. K+ ion channels (True/False)
- Distal tubular reabsorption is very crucial to reabsorb vital nutrients that are eliminated from body (True/False)
- 7. Drugs that are bioequivalent can be switched with one another in time of need (True/False)
- 8. Reason of various stages of metabolism is that, hydrophobic drugs cannot be efficiently eliminated by kidneys (True/False)
- 9. In the anesthesia procedure, flumazenil is mostly used in premedication (True/False)
- 10. Carbamazepine as an inducing agent increase the biotransformation of drug that ultimately lead to low level of plasma concentration (True/False)
- 11. According to therapeutic uses, lignocaine can be used both as surface and injectable anesthetic (True/False)
- 12. Benzodiazepines and barbiturates illicit the response after binding between an $\alpha 1$ and the $\gamma 2$ subunit in GABA receptor (True/False)
- Mostly phenytoin and sometimes Valproic acid exert its effects via inhibiting Na+ channels into rapidly firing neurons (True/False)
- 14. Status epilepticus the most serious, life-threatening condition characterized by convulsion and is generalized form of epilepsy (True/False)
- 15. Local anesthetics block the nerve impulse by blocking the ion channels at nerve terminalsi.e. K+ channels (True/False)