

Name:-

Shayiq-ur-Rahman.

ID:-

6952.

Assignment:-

Maxillofacial Surgery.

Program:-

BS (DT)

Submitted to:-

M-Natasha Kamran.

Q 1

a ORAL MUCOSA:-

- Oral mucosa is membrane lining inside of the mouth
- Stratified squamous epithelium
 - Lamina propria.

b Two Components:-

- Oral epithelium
- Lamina propria

ORAL MUCOSA:-

Masticatory.

Lining Mucosa

Specialized Mucosa.

c Connective tissue papilla:-

Irregular and upward projection of connective tissue.

Basement Membrane:-

Structureless layer about 2 micrometer thick.

Rete Ridges:-

or rete pegs that interdigitate with the connective tissue papilla

① Because in specialized mucosa
Lest Bed is present
therefore it is called specialized
mucosa.

Q2:

A Layer of Keratinized Surface epithelium:

Stratum Corneum.



Stratum Granulosum.



Stratum Spinosum.



Stratum Basale.

b Comprises layers of Germinativum.

Stratum Spinosum



Stratum Basale

C Pyknotic is the shrinkage and Condensation of cell.

Pyknotic nucleus found adrenal gland they are also found in Keratinocytes.

Q3

a Extraction:-

Tooth Extraction is the removal of a tooth from its socket in the bone.

Types

Simple extraction:-

Simple extraction is used to remove teeth that can be seen and are easily accessible.

Surgical Extractions

is that type of Extraction which typically require incision into Connective Tissue

b Type 2 Extraction

Surgical Extraction

C Indications:-

- Impacted tooth
- Severe periodontic disease

Contraindications:-

- periradicular Infection
- Space dentition-

Q4

a
Curettage

is the use of a curet to
Remove the soft tissue
Curettage is important after
Extraction because for proper
healing

b

The stitches will remove after
extraction in seven days.

- c
- 1) Gather the material Need it sharp
Scissors
 - 2) Sterilize the material
 - 3) wash the suture site.
 - 4) Stop if start bleeding.
 - 5) clean the area
 - 6) protect the wound.

Post operative Instruction will give to patients

Q6

After Extraction should be placed be the pressure on the gauze and covering the extraction site. ✓

if bleeding apply new gauze.

After surgery place a Cold Compress on face near the extraction site for 20 min.

Do not eat Drink hot food.

Do not use Straw.

Do not Spit.

Do not brush your teeth.

eat soft food.

Do not bite your lips.

b

Because the patient follow the specific instruction in order to remain comfortable and heal quickly.